

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## HEATING VALUES OF NATURAL GASES OF THE MIDCONTINENTAL FIELD.

By H. C. ALLEN.

WHILE a number of new gas pools hove been opened up in the midcontinental field in the last few years, no particular work of a chemical nature has been done on any of these gases since the work of Cady and McFarland.¹ Results of a preliminary examination of a number of these new gases are given in the following, together with some analyses of gases from the older pools of the field.

The thing of practical importance to the ordinary consumer of natural gas is its heating power, and these gases are of great interest in this line from the fact that they vary over such a wide range. This is illustrated in the following table:

TABLE I.

| No. | Location.<br>(Sec., twp., range.)  | CO <sub>2</sub> . | ннс. | Ο,.  | CH.   | С2Н". | Res.   | B.t.u. |
|-----|------------------------------------|-------------------|------|------|-------|-------|--------|--------|
| 1*  | Augusta, Kan. (500 feet)           | 0.13              | 0.00 | 0.00 | 10.54 | 1.64  | 87.69  | 129    |
| 2   | Cowley county, Kan. (24-33-6)      | 0.13              | 0.00 | 0.37 | 16.27 | 3.01  | 80.23  | 209    |
| 3   | Chautauqua county, Kan. (35-33-10) | 0.30              | 0.00 | 0.18 | 42.38 | 1.85  | 55.29  | 441    |
| 4   | Chautauqua county, Kan. (25-33-10) | 0.00              | 0.00 | 0.40 | 49.01 | 3.89  | 46.67  | 541    |
| 5*  | Ellsworth, Kan                     | 0.30              | 0.23 | 0.09 | 61.09 | 1.09  | 37.20  | 609    |
| 6*  | Ponca City, Okla. (4-25-2E)        | 0.05              | 0.00 | 0.40 | 44.60 | 14.86 | 40.10  | 688    |
| 7   | Kay county, Okla                   | 0.21              | 0.32 | 0.00 | 57.91 | 9.89  | -31.65 | 735    |
| 8*  | Chautauqua county, Kan. (15-34-13) | 0.55              | 0.00 | 0.81 | 85.53 | 0.15  | 12.95  | 839    |
| 9*  | Chautauqua county, Kan. (15-34-13) | 0.11              | 0.59 | 0.98 | 79.13 | 7.79  | 11.39  | 894    |
| 10  | Butler county, Kan. (2-28-4)       | 0.00              | 0.71 | 0.14 | 62.15 | 18.38 | 18.64  | 930    |
| 11  | Montgomery county, Kan             | 0.24              | 0.24 | 0.25 | 83.04 | 8.54  | 7.95   | 970    |
| 12  | Blackwell, Okla. (13-28-1W)        | 0.00              | 1.18 | 0.16 | 70.69 | 18.65 | 9.32   | 1025   |
| 13  | Cushing, Okla. (28-18-7)           | tr.               | 0.12 | tr.  | 70.74 | 21.64 | 7.49   | 1059   |
| 14  | Bartlesville, Okla. (12-26-14)     | 0.24              | 1.21 | 0.24 | 70.50 | 24.60 | 3.21   | 1125   |

All B.t.u. above are calculated to 0°C. and 760 mm. pressure from the following values: Heavy hydrocarbons (HHC), 1627; CH<sub>4</sub>, 966; C<sub>2</sub>H<sub>6</sub>, 1728.

\* Analyzed by Mr. E. E. Lyder. No. 8 from 530 feet of sand; No. 9 from 580 feet of sand.

<sup>1.</sup> University of Kansas Geol. Survey, vol. 9.

So far as the writer is aware, such a wide variation in composition and heating value is not found in the other natural-gas fields, and it has been a rather common practice to consider natural gases, with a few unimportant exceptions, to be of about the same composition and heating value.

The economic problem involved in the sale of these gases is a very material one, since some of these gases of low heating value are found in considerable quantities and the lower ones are practically valueless as fuels. While no effort is being made to sell the gas from the shallow Augusta or Dexter fields, gases of all the other grades listed above are being sold. The loss to the consumer in heat units, and consequently in money, from the use of a 500 B. t. u. gas amounts to 50 per cent when compared to the present gas supplied to Lawrence as a standard.

The following are typical analyses of gas from wells located in various parts of the midcontinental field:

| Location.<br>(Sec., twp., range.) | No. of<br>samples. | CO <sub>2</sub> . | ннс. | O <sub>2</sub> . | CH₄.  | С <sub>2</sub> Н <sub>6</sub> . | Res.  | B.t.u. |
|-----------------------------------|--------------------|-------------------|------|------------------|-------|---------------------------------|-------|--------|
| Pawhuska, (3-25-9)                | 3                  | 0.77              | 0.15 | 0.62             | 90.9  | 5.4                             | 2.4   | 971    |
| Cushing, (28-18-7)                | 5                  | tr.               | 0.12 | tr.              | 70.74 | 21.64                           | 7.49  | 1059   |
| Cushing, (6-18-7)                 | 2                  | 0.83              | 0.42 | 0.42             | 67.26 | 18.96                           | 12.14 | 984    |
| Bigheart, (Lot 298)               | 3                  | 0.86              | 0.15 | 0.30             | 78.90 | 16.43                           | 3.34  | 1048   |
| Hominy creek, (11-22-10)          | 2                  | 0.16              | 0.46 | 1.07             | 79.09 | 11.47                           | 7.76  | 970    |
| Hominy, (9-23-8)                  | 2                  | tr.               | 0.00 | tr.              | 75.65 | 8.69                            | 15.52 | 881    |
| Bartlesville, (12-26-14)          | 7                  | 0.13              | 1.03 | 0.39             | 72.02 | 21.16                           | 5.29  | 1078   |
| Hogshooter, (30-26-15)            | 5                  | 0.14              | 0.83 | 0.10             | 73.07 | 20.63                           | 5.23  | 1075   |
| Hogshooter, (33-26-14)            | 2                  | 1.55              | 0.00 | 0.39             | 91.82 | 2.35                            | 3.98  | 927    |
| North Osage, (2-28-10)            | 8                  | 0.40              | 0.80 | 0.54             | 72.64 | 13.62                           | 11.88 | 950    |
| Wann, (34-28-14)                  | 4                  | 0.48              | 0.49 | 0.32             | 76.66 | 16.26                           | 5.80  | 1029   |
| Wagner county, (6-19-15)          | 1                  | 1.60              | 1.53 | 0.36             | 69.01 | 17.31                           | 10.17 | 991    |
| Blackwell, (24-28-1W)             | 14                 | 0.00              | 0.32 | 0.47             | 71.40 | 20.13                           | 7.76  | 1043   |
| Blackwell,* (16-29-1E)            | 1                  | 0.05              | 0.10 | 0.11             | 60.54 | 6.22                            | 32.97 | 690    |
| Ponca,* (9-27-3E)                 | 5                  | 1.00              | 0.23 | 0.00             | 84.16 | 7.53                            | 7.08  | 947    |
| Ponca,* (4-25-2E)                 | 1                  | 0.05              | 0.00 | 0.40             | 44.60 | 14.86                           | 40.10 | 688    |

TABLE II. Oklahoma.

<sup>\*</sup> Analyses by Mr. E. E. Lyder.

TABLE III. Kansas.

| Location.<br>(Sec., twp., range.) | No. of samples. | CO <sub>2</sub> . | ннс. | O <sub>2</sub> . | CH <sub>4</sub> . | C2H6. | Res.  | B.t.u.   |
|-----------------------------------|-----------------|-------------------|------|------------------|-------------------|-------|-------|----------|
| Augusta, (a)                      | 10              |                   |      |                  |                   |       |       |          |
| Sedan, (b)                        | 13              |                   |      |                  |                   |       |       | <b>.</b> |
| Sedan, (c)                        | 6               |                   |      |                  | <b>.</b>          |       |       |          |
| Arkansas City,* (15-35-2)         | 2               | 0.00              | 0.38 | 0.10             | 67.36             | 14.68 | 17.46 | 908      |
| Arkansas City,* (18-34-4)         | 1               | 0.00              | 0.43 | 0.00             | 73.28             | 17.10 | 9.18  | 1006     |
| Allen county, (9-26-20)           | 1               | 0.35              | 1.15 | 0.39             | 79.80             | 4.33  | 14.00 | 865      |
| Winfield,* (12-33-4)              | 5               | 0.36              | 0.00 | 0.24             | 77.90             | 9.51  | 11.98 | 912      |
| Havana                            | 2               | 0.00              | 0.45 | 0.20             | 72.56             | 13.46 | 13.34 | 955      |
| Benedict, (16-28-15)              | 1               | 0.12              | 0.00 | 0.46             | 92.30             |       | 4.30  | 900      |

<sup>\*</sup> Analyses by Mr. E. E. Lyder.
(a) See Table I, No. 1, for 1400 feet of gas.

TABLE IV. City Supplies.

| Сіту.              | ]    | Date.    | CO2. | ннс. | 02.  | CH₄.  | C2H6. | Res.  | B.t.u. |
|--------------------|------|----------|------|------|------|-------|-------|-------|--------|
| Lawrence, Kan      | Aug. | 19, 1913 | 0.00 | 0.23 | 0.35 | 88.18 | 4.90  | 6.12  | 941    |
| Lawrence, Kan      | Jan. | 9, 1914  | 0.62 | 0.62 | 0.17 | 82.38 | 9.92  | 6.22  | 977    |
| Lawrence, Kan      | Feb. | 23, 1914 | 0.72 | 0.57 | tr.  | 86.42 | 7.23  | 5.06  | 969    |
| Lawrence, Kan      | Nov. | 24, 1914 | 1.28 | 1.55 | 0.93 | 79.00 | 7.28  | 10.00 | 913    |
| Lawrence, Kan      | Jan. | 15, 1915 | 0.71 | 0.24 | 0.08 | 85.77 | 7.97  | 5.25  | 970    |
| Lawrence, Kan.*    | Mar. | 14, 1915 | 0.79 | 0.13 | 0.35 | 81.75 | 11.14 | 5.82  | 979    |
| Lawrence, Kan.*    | Apr. | 15, 1915 | 0.13 | 0.54 | 0.29 | 88.30 | 6.82  | 3.82  | 975    |
| Udall, Kan         | Dec. | 18, 1913 | 0.27 | 0.37 | 0.44 | 63.00 | 5.67  | 30.35 | 713    |
| Udall, Kan         | Feb. | 18, 1914 | 0.00 | 0.00 | 0.38 | 73.39 | 17.73 | 8.51  | 1015   |
| Wellington, Kan    | Dec. | 18, 1913 | 0.20 | 0.12 | 0.36 | 65.18 | 10.20 | 23.96 | 808    |
| Wellington, Kan    | Feb. | 18, 1914 | 0.00 | 0.60 | 0.31 | 75.53 | 15.42 | 8.15  | 1005   |
| Winfield, Kan      | Feb. | 18, 1914 | 0.00 | 0.47 | 0.63 | 73.82 | 16.87 | 8.21  | 1013   |
| Burden, Kan        | Feb. | 18, 1914 | 0.93 | 0.00 | 0.31 | 79.26 | 10.33 | 9.17  | 944    |
| Augusta, Kan       | Feb. | 21, 1914 | 0.00 | 1.01 | tr.  | 75.68 | 10.97 | 12.33 | 956    |
| Ottawa, Kan        | Feb. | 21, 1915 | 0.40 | 0.53 | 0.54 | 88.93 | 4.68  | 4.93  | 948    |
| Wichita, Kan       | Dec. | 18, 1913 | 0.27 | 0.76 | 0.22 | 59.88 | 15.69 | 23.13 | 862    |
| Wichita, Kan       | Feb. | 10, 1914 | 0.36 | 0.72 | 0.24 | 64.71 | 17.66 | 16.28 | 942    |
| Hutchinson, Kan    | Feb. | 10, 1914 | 0.46 | 0.70 | 0.23 | 66.20 | 17.24 | 15.17 | 949    |
| Arkansas City, Kan | Dec. | 18, 1913 | 0.16 | 0.17 | 0.20 | 52.06 | 33.42 | 15.10 | 1066   |
| El Dorado, Kan     | Dec. | 8, 1913  | 0.53 | 0.00 | 0.59 | 45.70 | 30.81 | 22.49 | 974    |
| Fort Worth, Tex    | Jan. | 25, 1914 | 0.00 | 0.63 | tr.  | 46.40 | 13.96 | 39.01 | 700    |
| Dallas, Tex        | Jan. | 24, 1914 | 0.00 | 0.51 | 0.25 | 40.92 | 17.96 | 40.37 | 714    |

<sup>\*</sup>Analyses by Mr. E. E. Lyder.

The above work is preliminary, in a way, to a general survey of the midcontinental field that is being made by the University of Kansas, and which is to be published later.

<sup>(</sup>b) See Table I, No. 4. (c) See Table I, No. 3.